

# Physiotherapy in cystic fibrosis

## A comprehensive clinical overview

**Arietta Spinou MSc, PhD**

Health, Sports and Bioscience,  
University of East London, UK

**Key words:**

- Physiotherapy
- Cystic Fibrosis
- Airway Clearance
- Chest Physiotherapy
- Exercise

**Correspondence:**

Arietta Spinou,  
Lecturer in Physiotherapy,  
Health, Sports and Bioscience,  
University of East London, Water Lane,  
Stratford, E15 4LZ, London, UK  
E-mail: a.spinou@uel.ac.uk

### SUMMARY

Physiotherapy remains the cornerstone of cystic fibrosis (CF) management alongside medical treatment. Traditionally, physiotherapy intervention focussed on airway clearance during the clinically stable stage and chest infections. Research evidence consistently supports greater mucus clearance with chest physiotherapy compared to cough alone or no treatment. Various methods and techniques of airway clearance have been developed and investigated, and data suggest that most of them are of similar effectiveness. Nowadays physiotherapy management also extends to other areas, supported by studies and clinical practice. The physiotherapists plan, supervise and follow-up systematic exercise or personalised rehabilitation programs, which, similarly to airway clearance, are recommended in all patients with CF. Furthermore, based on a comprehensive assessment, physiotherapists incorporate the management of accompanying musculoskeletal problems such as back pain and postural disorders, as well as urine incontinence issues. In the era that aims to improve quality of life, it is essential that physiotherapists are aware of specific conditions that might affect the management of CF. Their role is to work alongside and within the CF multi-disciplinary team throughout patient's treatment and consistently support the patient and carers, in particular whilst on clinical pathways of the lung transplantation and palliative care.

*Pneumon 2018, 31(1):35-43.*

### INTRODUCTION

Cystic fibrosis (CF) is a recessive genetic disease that affects the patient on multiple systems, with profound manifestations in the respiratory and digestive systems<sup>1</sup>. It is characterised by the mutation and therefore dysfunction of the gene for the cystic fibrosis transmembrane conductance regulator (CFTR). This protein mainly functions as an ion channel, regulating fluid volume on epithelial surfaces via chlorine secretion and inhibition of sodium resorption. In the airways of the patients with CF, dysfunction

of the CFTR results in periciliary liquid layer depletion<sup>2</sup>. Clinically, patients with CF present abnormal consistency and high volumes of sputum, cough, dyspnoea, bronchiectasis and weight loss. As the survival of these patients is increasing, it is crucial that health care professionals address symptoms and support individuals in evolving issues developed throughout their life span.

Physiotherapy is an integral part of the therapeutic management of CF patients, both at the clinically stable stage of the disease and during respiratory infections. In the past, physiotherapy was focused on airway clearance, also known as chest physiotherapy, by teaching or applying methods such as the postural drainage with or without the additional application of manual techniques<sup>3</sup>. Postural drainage of the tracheobronchial tree uses specific gravitational positions to assist mucus mobilisation downwards (towards the mouth) within the airways. Manual techniques (percussions, vibrations and/or shakes) use mechanical forces to assist the detachment of mucus from the airway epithelium and its mobilisation. Nowadays, the choice of airway clearance techniques has been expanded to methods such as the autogenic drainage, the active cycle of breathing techniques (ACBT), the use of positive expiratory pressure (PEP) devices with or without oscillation, and others. Still, modern physiotherapy in CF also includes the assessment of the cardiovascular system and improvement of the patient's fitness level, muscle strength and endurance through exercise, as well as specialised interventions to improve musculoskeletal symptoms of pain, posture and incontinence<sup>4</sup>.

## PHYSIOTHERAPY

### Airway clearance

Patient education, application and monitoring of the airway clearance techniques remain the main physiotherapy treatment for patients with CF<sup>4</sup>. Physiotherapists facilitate the establishment of an individualised airway clearance routine by supporting patients and their families to establish regular regimes during a clinically stable stage and have an escalation plan for disease exacerbations<sup>5</sup>. Airway clearance is usually performed on a daily basis and as required. *The selected method applied, duration and frequency of each session are tailored to the patient, their general health condition and the severity of the disease.* For instance, airway clearance becomes more regular during exacerbations or hospitalisations<sup>6</sup>. Hospitalisations also provide an opportunity for physiotherapists to re-assess

the effectiveness of daily airway clearance and provide appropriate feedback and guidance for improving the patient's usual technique prior to discharge.

Table 1 presents the main categories of airway clearance techniques and methods in CF. These can be used in isolation or in combination regimes. Assessment of effectiveness is based on measuring sputum volume or weight, lung function by spirometry, frequency of hospitalisations and quality of life. Airway clearance is extensively supported in the literature when compared to no airway clearance or cough alone<sup>4,7-9</sup>. A recent systematic review supported a significant increase in the amount of sputum (wet or dry) in the patient groups that applied airway clearance using postural drainage with or without the addition of manual techniques or using PEP, compared to spontaneous cough or not using any technique<sup>7</sup>. The weight of the sputum was higher after the application of the active cycle of breathing techniques compared to the use of the flutter (an oscillating PEP device) or high frequency chest wall oscillation (vest)<sup>10</sup>. The weight of the sputum expectorated was greater after using the PEP mask compared to autogenic drainage, postural drainage positions and their combination, although this difference was short-term (up to one week)<sup>11</sup>. On the other hand, there was no difference in the amount of the expectorated mucus after autogenic drainage compared to the flutter, or between the high frequency chest wall oscillation compared to the autogenic drainage or the PEP mask for longer time-intervals<sup>10,12</sup>.

Systematic reviews did not show significant differences in the lung function (FEV<sub>1</sub>) of adult patients fol-

**TABLE 1.** Common airway clearance techniques and methods.

#### Airway clearance techniques

- Postural drainage
- Manual techniques
- Active cycle of breathing techniques (ACBT)
- Autogenous drainage (AD)
- Positive expiratory pressure (PEP) devices (PEP mask, Pari-PEP, etc)
- Positive expiratory pressure (PEP) devices with oscillation (flutter, acapella, cornet, etc.)
- Intermittent Positive Pressure Breathing (IPPB)
- High frequency chest wall oscillation (HFCWO) or vest
- Non-invasive mechanical ventilation (NIV)
- Aerobic exercise

lowing the use of PEP, when assessed patients prior and immediately after a physiotherapy session or up to 3 months later<sup>7,10,11,13</sup>. Additionally, the lung function did not change after applying the active cycle of breathing techniques in combination with the PEP mask, postural drainage with or without manual techniques, or the high frequency chest wall oscillation<sup>12</sup>. However, treatment in children and adolescents that was applied up to one year showed 6% increase in FEV<sub>1</sub> with the use of PEP<sup>13</sup>.

Regarding the hospitalisation frequency, no differences were found for those who practiced the active cycle of breathing techniques compared to the postural drainage with or without manual techniques<sup>12</sup>. The number of hospitalisations, however, was lower for those who used PEP than the patients who used the flutter (5 vs 18 hospitalisations, respectively)<sup>10</sup>. Similarly, fewer patients used intravenous antibiotics from the group that used PEP devices, compared to the group of the high frequency chest wall oscillation<sup>13</sup>.

For the quality of life, there is no difference amongst techniques and devices, such as the postural drainage with or without manual techniques, active cycle of breathing techniques, autogenic drainage, PEP mask, flutter, and cornet<sup>10,12,13</sup>. However, patients preferred the PEP mask for long-term use (>1 month), and also preferred seating instead of using postural drainage positions<sup>10,11,13</sup>. Autogenic drainage was preferred among children between 12-18 years old, compared to postural drainage in combination with manual techniques<sup>14</sup>.

Important factors for the success of the selected airway clearance plan are the compliance to treatment and patient satisfaction. Factors that increase the rate of compliance are good patient knowledge of the technique and confidence in its application, independence and preference<sup>15,16</sup>. Evidence indicate that patients who receive help, those who produce more sputum, and children with CF whose parents believe in the necessity of treatment are those with higher compliance in airway clearance<sup>17,18</sup>.

## Airway clearance adaptations

### *Mucolytics and other agents*

Patients with CF often receive medications that aim to increase the effectiveness of airway clearance, such as nebulised hypertonic saline (3% to 7% NaCl), dornase alpha (DNase), and mannitol. The use of inhaled hypertonic saline (osmotic pressure >0.9% NaCl) in patients with CF is considered to improve the rheological characteristics of sputum and increase the hydration of the airway epithelium; thus, increase the sputum motility and facilitate the mucus clearance<sup>19</sup>. There is good evidence that the use of hypertonic saline reduces the incidence of respiratory infections, increases FEV<sub>1</sub>, and improves the quality of life, although the changes are not maintained in the long term (48 weeks)<sup>20,21</sup>. During the hospitalisation of patients with CF, hypertonic saline improves the chances of quick return of the lung function (FEV<sub>1</sub>) to pre-infectious levels<sup>22</sup>. With regards to timing the hypertonic saline administration, a recent systematic review supports its use before or during the performance of airway clearance, rather than its administration afterwards<sup>23</sup>. If the prescribed doses are two, it is recommended to administer one in the morning and one in the evening, and if the patient receives a single dose this is given at a convenient time chosen by the patient<sup>23</sup>.

Dornase alpha (DNase) is a recombinant human deoxyribonuclease that reduces sputum viscosity by selectively hydrolysing the large extracellular DNA molecules contained in the mucus into smaller structures, thereby increasing the potential for its elimination<sup>24</sup>. This drug is administered via a jet-nebuliser device and has been shown to reduce the incidence of respiratory infections, increase respiratory function, and improve quality of life<sup>24</sup>. With regards to timing its administration, it appears that using DNase before or after airway clearance does not have any difference in improving lung function (FEV<sub>1</sub> and FVC) or patient's quality of life<sup>25,26</sup>. In clinical practice, physiotherapy often follows the proposed guidelines of the pharmaceutical company to perform airway clearance 30 minutes after the DNase administration<sup>27</sup>.

Inhaled mannitol is a naturally occurring sugar alcohol which enhances osmosis, causing mucus hydration<sup>28</sup>. Inhaled mannitol is administered as dry powder (capsules) using an inhaler. As demonstrated by two 26-week multicentre studies with a total number of 600 participants with CF, inhaled mannitol improves the respiratory function of patients but does not improve their quality of life<sup>29,30</sup>. Although its use usually precedes airway clearance in clinical practice, there is no research data to compare different timings of administration.

Inhaled mannitol is a naturally occurring sugar alcohol which enhances osmosis, causing mucus hydration<sup>28</sup>. Inhaled mannitol is administered as dry powder (capsules) using an inhaler. As demonstrated by two 26-week multicentre studies with a total number of 600 participants with CF, inhaled mannitol improves the respiratory function of patients but does not improve their quality of life<sup>29,30</sup>. Although its use usually precedes airway clearance in clinical practice, there is no research data to compare different timings of administration.

### *Haemoptysis*

Haemoptysis is a major change in the patient's clinical presentation and may be life-threatening. The physiotherapy assessment should include questions about sputum description and reference to current or past haemoptysis episodes. Active frank haemoptysis (>100-1000 ml haemoptysis in 24 hours or 48 hours) is treated

exclusively medically, e.g. with bronchial embolisation of the arteries or thoracic surgery, while the airway clearance treatment is temporarily discontinued<sup>31,32</sup>. In moderate or low haemoptysis, physiotherapists, in collaboration with the medical team, decide whether or not it is appropriate to continue airway clearance using clinically reasoning. If the treatment is appropriate and safe to continue, then the active cycle of breathing techniques or autogenic drainage is often selected over other techniques.

### ***Pneumothorax***

Spontaneous pneumothorax is a common complication in patients with CF. It is associated with a reduction in pulmonary function and 50-90% chance of recurrence<sup>32,33</sup>. If the pneumothorax occurs for the first time and it is small, then it can be treated conservatively with oxygen supply<sup>34</sup>. In patients continuing airway clearance, it is suggested to liaise with the medical team for adding humidification to the oxygen supply and ensuring adequate analgesia for the duration of the treatment sessions<sup>35</sup>. In the case of large pneumothorax (>2 cm between parietal pleura and visceral pleura) or recurrent pneumothorax, chest drainage is performed using thoracic catheters, while patients might get pleurodesis in resistant cases<sup>34</sup>. Positive pressure devices such as PEP, flutter and acapella are contraindicated in the presence of pneumothorax<sup>34</sup>. Regarding physical activity, patients need to be engaged with moderate activities but should avoid bearing weights over 2 kg or strenuous aerobic exercise for a period of two to six weeks after the complete drainage of the pneumothorax<sup>34</sup>.

### **Exercise**

Exercise is an integral part of the comprehensive physiotherapy intervention for patients with CF<sup>36</sup>. American College of Sports Medicine guidelines advocate 3-5 sessions of moderate exercise per week, with the aim to adopt exercise as a way of living<sup>37</sup>. Benefits of specific exercise modalities in cystic fibrosis are yet to be identified in methodologically strong studies<sup>38</sup>. Despite research interest, evidence has not established the effectiveness of inspiratory muscle training on this group of patients, therefore this is currently not routinely incorporated in the CF treatment. In the clinical setting, the assessment of patients with CF uses simple and cost-effective exercise field tests, such as the 6-minute walk test (6MWT) and the incremental shuttle walk test (ISWT), whilst the level of dyspnoea is assessed using the Borg dyspnoea scale<sup>39</sup>.

Exercise can theoretically assist airway clearance through the kinetic forces and vibrations generated within the airways, but it cannot substitute for the formal airway clearance<sup>40</sup>. When compared to airway clearance techniques, moderate aerobic exercise leads to less mucus expectoration<sup>41</sup>. Also, exercise as a single agent does not increase cough immediately after its completion, although it improves the subjective ease of sputum clearance<sup>42</sup>. Clinically, exercise is mainly used additionally to airway clearance, as a means to improve the exercise capacity of the patient and is usually performed before the implementation of airway clearance.

### **Exercise considerations**

#### ***Musculoskeletal and postural issues***

Back and thoracic pain are frequently reported in patients with CF, although they do not have an effect on lung function<sup>43,44</sup>. Higher thoracic kyphosis is associated with lower lung function, but nowadays it is more uncommon compared to a few years ago<sup>45</sup>. Low bone density and osteopenia is also a common issue in patients with CF<sup>46,47</sup>. Counselling and appropriate exercise programs from physiotherapists can potentially address and improve these postural and structural issues<sup>36</sup>.

#### ***Urinary incontinence***

Surveys show that urinary incontinence in patients with CF is reported in 30% to 68% of women or girls and 5% to 16% of men or boys<sup>48-51</sup>. The dynamic pressure created during coughing is potentially a key mechanism of CF urinary incontinence, although it may not be the only one<sup>52</sup>. Coughing, sneezing, laughing and spirometry are among the activities that trigger urinary incontinence incidents<sup>53</sup>. Incontinence worsens during respiratory infections and has been associated with poorer quality of life and higher anxiety and depression scores<sup>51,54,55</sup>. Assessing incontinence using screening tools and clarifying questions should be an integral part of the CF physiotherapy assessment, regardless of gender<sup>56</sup>. Physiotherapy treatment of urinary incontinence includes counselling and specialised training involving pelvic floor exercises, such as Kegel exercises<sup>55,57,58</sup>.

#### ***Diabetes mellitus***

Diabetes mellitus is associated with CF and is the most common comorbidity of the disease, occurring in up to 20-50% of adult patients<sup>59-61</sup>. This comorbidity requires the co-operation of the physiotherapists with the

endocrine team, especially for the patients who require insulin therapy<sup>62</sup>. Additionally, the presence of diabetes mellitus needs to be considered in the physiotherapy plan, mainly in the exercise prescription and performance. In this case, the proper scheduling of the meal times or insulin intake is essential.

## Quality of life

Over time and as the CF severity and symptoms progress, the quality of life of patients is deteriorating. Females with CF often report poorer quality of life compared to their male age-matched peers<sup>63</sup>. Although the correlation between lung function and quality of life is weak to moderate, patients with better lung function report higher quality of life<sup>54</sup>. Also, the presence of *Pseudomonas aeruginosa* and frequent respiratory infections appear to have a negative impact on the quality of life of patients<sup>54</sup>.

Researchers and clinicians can use a number of validated questionnaires for the assessment of quality of life in people with CF. Those include: generic questionnaires or questionnaires for a specific disease symptom, such as the Short Form-36 (SF-36) and the Leicester Cough Questionnaire, respectively<sup>64,65</sup>; disease-specific questionnaires, such as the Manchester Questionnaire, the Cystic Fibrosis Questionnaire-Revised and the Cystic Fibrosis-Quality of Life<sup>64,66-69</sup>; and questionnaires for babies and children of young age, such as the Modified Parent Cystic Fibrosis Questionnaire-Revised<sup>70</sup>.

## Special considerations

### *Long term oxygen therapy and non-invasive ventilation*

A recent systematic review in patient with CF did not show long-term benefits from the long-term oxygen therapy, in survival, respiratory function or cardiovascular health, although it showed improved school or work attendance rates<sup>71</sup>. When oxygen is administered during exercise only, it helps to improve oxygenation, reduces the feeling of dyspnoea and increases the duration of the exercise<sup>71,72</sup>. However, supplemental oxygen during exercise in patients with initially low arterial oxygen values appears to cause hypercapnia in the short term (PCO<sub>2</sub> up to 16 mmHg)<sup>71</sup>. Also, oxygen therapy during sleep improves oxygenation, but is accompanied by small hypercapnia<sup>71</sup>. The use of supplemental oxygen should follow the established clinical guidelines that are based on hypoxia (PaO<sub>2</sub> ≤55 mmHg or 60 mmHg) and the presence of clinical symptoms<sup>73</sup>.

Non-invasive ventilation (NIV) is used in patients with

CF on respiratory failure, hypoventilation during sleep, as well as a bridge to lung transplantation<sup>3</sup>. For patients with severe clinical presentation where airway clearance causes fatigue and high levels of dyspnoea, NIV can be used to assist airway clearance<sup>74</sup>. The use of NIV during the physiotherapy session facilitates mucus expectoration and reduces the sensation of dyspnoea during the treatment compared to other techniques particularly for patients with low lung function<sup>75</sup>. However, the long-term effects of NIV on airway clearance need further investigation<sup>76</sup>.

## *Paediatric population*

Choosing a treatment plan for children with CF is based on age, clinical presentation and certain social criteria<sup>77</sup>. There is no agreement on the most appropriate starting age for airway clearance. A proposal for early disease management (pre-symptomatic) is to carefully monitor the clinical presentation of children and adopt an active treatment plan following the onset of symptoms<sup>78</sup>. At young ages, where the child can not follow instructions and cooperate, assisted autogenic drainage or PEP devices with a child mask can be used. Physiotherapists are also responsible for educating the child's parents or carers for appropriate evaluation of the child's symptoms and treatment implementation as required<sup>79</sup>. Postural drainage with tilt (head-down positions) is no longer advised for babies, as it has been shown to increase the gastroesophageal reflux<sup>80</sup>.

As children grow older, they can more actively participate in their treatment. Children over 3 years old can also use an airway clearance game, the bubble PEP. This is a positive-pressure home-made device, where children are encouraged to generate soap bubbles by breathing out through a small plastic tube and into a bottle of soapy water<sup>81</sup>. According to the UK Cystic Fibrosis Foundation, at the age of 6 years or more, the use of nebulised hypertonic saline can be initiated in combination with airway clearance<sup>82</sup>. Also, at all ages, activity games and engagement with exercise are encouraged and used, for instance racing, trampolines and exercises using a gym ball<sup>83</sup>.

## *Palliative care*

CF is a disease that limits life expectancy and requires discipline and consistency to many hours of daily treatment. As a result, its psychological impact should not be ignored<sup>84</sup>. If patients are in respiratory failure and in lung transplantation list, pulmonary rehabilitation is the treatment priority, alongside the aim to relieve symptoms. Working in line with the patient's wishes is very impor-

tant, particularly during the palliative care stage. Airway clearance of less active patient participation (eg. postural drainage), massage and some dyspnoea relieving positions could be applied during this stage, if they provide comfort to the patient<sup>85</sup>.

## CONCLUSIONS

CF management is highly demanding, mainly aiming to the reduction and treatment of chest infections, improvement of quality of life and increase of life expectancy. Physiotherapy is an integral part of the patient's daily treatment routine, and additionally to airway clearance other important issues should be addressed. International clinical guidelines suggest access to specialised physiotherapy care both during a clinically stable stage of the disease and during respiratory infections. At the clinically stable stage, patients should be evaluated by physiotherapists every 3-6 months to re-evaluate and op-

timize their treatment plan. During respiratory infections, physiotherapy interventions are intensified according to the clinical presentation. Although in CF airway clearance is the cornerstone of physiotherapy treatment, physiotherapists work beyond the respiratory system and play an important role in the management of other issues, mainly using individualised exercise programmes. The exercise programmes need to be tailored to patient-related needs and issues, such as pain, diabetes and incontinence. This way, the patient-centred and individualised treatment follows the international standards and clinical guidelines.

## CONFLICT OF INTEREST DECLARATION

No conflict of interest.

## FUNDING

None.

---

## ΠΕΡΙΛΗΨΗ

### Φυσικοθεραπεία στην κυστική ίνωση: Μια περιεκτική κλινική ανασκόπηση

Αριέττα Σπίνου MSc, PhD

Λέκτορας Φυσικοθεραπείας, Health, Sports and Bioscience, University of East London, UK

*Η φυσικοθεραπεία παραμένει μια από τις κύριες μεθόδους διαχείρισης την κυστικής ίνωσης, σε συνδυασμό με την ιατρική θεραπεία. Παραδοσιακά, η φυσικοθεραπεία επικεντρώνονταν στον τραχειοβρογχικό καθαρισμό κατά τη διάρκεια της κλινικά σταθερής φάσης και των αναπνευστικών λοιμώξεων, με τα ερευνητικά δεδομένα να υποστηρίζουν την αποτελεσματικότητά της συγκριτικά με τον βήχα ή τη μη θεραπεία. Διάφορες μέθοδοι και τεχνικές τραχειοβρογχικού καθαρισμού έχουν αναπτυχθεί και διερευνηθεί, και τα δεδομένα προτείνουν ότι οι περισσότερες από αυτές είναι παρόμοιας αποτελεσματικότητας. Επιπλέον, σήμερα, οι έρευνες και η κλινική πρακτική επεκτείνουν τη φυσικοθεραπευτική διαχείριση πέραν του αμιγώς αναπνευστικού συστήματος. Οι φυσικοθεραπευτές σχεδιάζουν, επιβλέπουν και επανελέγχουν τη συστηματική άσκηση ή εξατομικευμένο πρόγραμμα αποκατάστασης, που ομοίως με τον τραχειοβρογχικό καθαρισμό συστήνεται σε όλους τους ασθενείς με κυστική ίνωση. Ακόμα, όταν χρειάζεται και με βάση μια ολοκληρωμένη αξιολόγηση, η φυσικοθεραπεία πραγματεύεται τη διαχείριση συνοδών μυοσκελετικών προβλημάτων όπως οσφυαλγίας, εργονομικών προβλημάτων στάσης και ακράτειας. Σε μια εποχή που στοχεύει στη βελτίωση της ποιότητας ζωής, οι φυσικοθεραπευτές είναι απαραίτητο να γνωρίζουν τις ειδικές περιπτώσεις που επηρεάζουν τη διαχείριση της κυστικής ίνωσης. Ο ρόλος τους είναι να εργάζονται σε συνεργασία με την πολύ-επιστημονική ομάδα για την υποστήριξη των ασθενών και του περιβάλλοντός τους, ιδιαίτερα όταν οι ασθενείς είναι σε αναμονή για μεταμόσχευση ή κατά την παρηγορητική φροντίδα.*

**Πνεύμων 2018, 31(1):35-43.**

---

**Λέξεις - Κλειδιά:** φυσικοθεραπεία, κυστική ίνωση, τραχειοβρογχικός καθαρισμός, αναπνευστική φυσικοθεραπεία, άσκηση

## REFERENCES

1. Elborn JS. Cystic fibrosis. *Lancet* 2016; 388:2519-31.
2. Matsui H, Grubb BR, Tarran R, et al. Evidence for periciliary liquid layer depletion, not abnormal ion composition, in the pathogenesis of cystic fibrosis airways disease. *Cell* 1998; 95:1005-15.
3. Pryor JA, Tannenbaum E, Scott SF, et al. Beyond postural drainage and percussion: Airway clearance in people with cystic fibrosis. *Journal of Cystic Fibrosis* 2010; 9:187-92.
4. Bott J, Blumenthal S, Buxton M, et al. Guidelines for the physiotherapy management of the adult, medical, spontaneously breathing patient. *Thorax* 2009; 64:(Suppl 1):i1-51.
5. National Guideline Alliance (UK). Cystic fibrosis: diagnosis and management. London: National Institute for Health and Care Excellence (UK); 2017 Oct.
6. Flume PA, Mogayzel JP Jr, Robinson AK, et al. Cystic fibrosis pulmonary Guidelines. Treatment of pulmonary exacerbations. *American Journal of Respiratory and Critical Care Medicine* 2009; 180:802-8.
7. Warnock L, Gates A. Chest physiotherapy compared to no chest physiotherapy for cystic fibrosis. *Cochrane Database Syst Rev* 2015: Cd001401.
8. Bradley JM, Moran FM, Elborn JS. Evidence for physical therapies (airway clearance and physical training) in cystic fibrosis: an overview of five Cochrane systematic reviews. *Respir Med* 2006; 100:191-201.
9. Main E, Prasad A, van der Schans CP. Conventional chest physiotherapy compared to other airway clearance techniques for cystic fibrosis. *Cochrane Database of Systematic Reviews*, 2005.
10. Morrison L, Agnew J. Oscillating devices for airway clearance in people with cystic fibrosis. *Cochrane Database of Systematic Reviews* 2014.
11. Elkins MR, Jones A, van der Schans C. Positive expiratory pressure physiotherapy for airway clearance in people with cystic fibrosis. *Cochrane Database Syst Rev* 2006: Cd003147.
12. McKoy NA, Wilson LM, Saldanha IJ, Odelola OA, Robinson KA. Active cycle of breathing technique for cystic fibrosis. *Cochrane Database of Systematic Reviews* 2016.
13. McIlwaine M, Button B, Dwan K. Positive expiratory pressure physiotherapy for airway clearance in people with cystic fibrosis. *Cochrane Database of Systematic Reviews* 2015.
14. McIlwaine M, Wong LT, Chilvers M, Davidson GF. Long-term comparative trial of two different physiotherapy techniques; postural drainage with percussion and autogenic drainage, in the treatment of cystic fibrosis. *Pediatr Pulmonol* 2010; 45:1064-9.
15. Lester MK, Flume PA. Airway-clearance therapy guidelines and implementation. *Respir Care* 2009; 54:733-50; discussion 751-3.
16. Homnick DN. Making airway clearance successful. *Paediatric Respiratory Reviews* 2007; 8:40-5.
17. Abbott J, Dodd M, Bilton D, Webb AK. Treatment compliance in adults with cystic fibrosis. *Thorax* 1994; 49:115-20.
18. Goodfellow NA, Goodfellow AN, Hawwa FA, Reid JMA, Horne R, Shields DM, McElroy CJ. Adherence to treatment in children and adolescents with cystic fibrosis: a cross-sectional, multi-method study investigating the influence of beliefs about treatment and parental depressive symptoms. *BMC Pulm Med* 2015; 15:43.
19. Donaldson SH, Bennett WD, Zeman KL, Knowles MR, Tarran R, Boucher RC. Mucus clearance and lung function in cystic fibrosis with hypertonic saline. *N Engl J Med* 2006; 354:241-50.
20. Wark P, McDonald VM. Nebulised hypertonic saline for cystic fibrosis. *Cochrane Database of Systematic Reviews* 2009.
21. Elkins MR, Robinson M, Rose RB, et al. A controlled trial of long-term inhaled hypertonic saline in patients with cystic fibrosis. *N Engl J Med* 2006; 354:229-40.
22. Dentice RL, Elkins RM, Middleton GP, et al. A randomised trial of hypertonic saline during hospitalisation for exacerbation of cystic fibrosis. *Thorax* 2016; 71:141-7.
23. Elkins M, Dentice R. Timing of hypertonic saline inhalation for cystic fibrosis. *Cochrane Database Syst Rev* 2016; 12: Cd008816.
24. Yang C, Chilvers M, Montgomery M, Nolan SJ. Dornase alfa for cystic fibrosis. *Cochrane Database of Systematic Reviews* 2016.
25. Dentice R, Elkins M. Timing of dornase alfa inhalation for cystic fibrosis. *Cochrane Database of Systematic Reviews*, 2016.
26. Bishop JR, Erskine OJ, Middleton PG. Timing of dornase alfa inhalation does not affect the efficacy of an airway clearance regimen in adults with cystic fibrosis: a randomised crossover trial. *J Physiother* 2011; 57:223-9.
27. Shak S, Capon DJ, Hellmiss R, Marsters SA, Baker CL. Recombinant human DNase I reduces the viscosity of cystic fibrosis sputum. *Proc Natl Acad Sci U S A*, 1990; 87:9188-92.
28. Daviskas E, Anderson SD, Jaques A, Charlton B. Inhaled mannitol improves the hydration and surface properties of sputum in patients with cystic fibrosis. *CHEST Journal* 2010; 137:861-8.
29. Nolan SJ, Thornton J, Murray CS, Dwyer T. Inhaled mannitol for cystic fibrosis. *Cochrane Database Syst Rev* 2015: Cd008649. doi: 10.1002/14651858.CD008649.pub2.
30. Bilton D, Bellon G, Charlton B, et al. Pooled analysis of two large randomised phase III inhaled mannitol studies in cystic fibrosis. *Journal of Cystic Fibrosis*, 2013; 12:367-76.
31. Ibrahim WH. Massive haemoptysis: the definition should be revised. *European Respiratory Journal* 2008; 32:1131-2.
32. Flume PA, Mogayzel PJ Jr, Robinson KA, Rosenblatt RL, Quitell L, Marshall BC; Clinical Practice Guidelines for Pulmonary Therapies Committee; Cystic Fibrosis Foundation Pulmonary Therapies Committee. Cystic fibrosis pulmonary guidelines: pulmonary complications: hemoptysis and pneumothorax. *Am J Respir Crit Care Med* 2010; 182:298-306.
33. Flume PA. Pneumothorax in cystic fibrosis. *Curr Opin Pulm Med* 2011; 17:220-5.
34. MacDuff A, Arnold A, Harvey J. Management of spontaneous pneumothorax: British Thoracic Society Pleural Disease Guideline 2010. *Thorax* 2010; 65:Suppl 2:ii18-31.
35. Flume PA. Pneumothorax in cystic fibrosis. *Chest* 2003; 123:217-21.
36. Smidt N, de Vet HC, Bouter LM, et al. Effectiveness of exercise

- therapy: a best-evidence summary of systematic reviews. *Aust J Physiother* 2005; 51:71-85.
37. Haskell WL, Lee IM, Pate RR, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007; 39:1423-34.
  38. Radtke T, Nolan SJ, Hebestreit H, Kriemler S. Physical exercise training for cystic fibrosis. *Cochrane Database Syst Rev* 2015;Cd002768. doi: 10.1002/14651858.CD002768.pub3.
  39. Holland AE, Spruit MA, Troosters T, et al. An official European Respiratory Society/American Thoracic Society technical standard: field walking tests in chronic respiratory disease. *Eur Respir J* 2014; 44:1428-46.
  40. McCool FD, Rosen MJ. Nonpharmacologic airway clearance therapies: ACCP evidence-based clinical practice guidelines. *Chest* 2006; 129(1 Suppl):250s-259s.
  41. Bilton D, Dodd ME, Abbot JV, Webb AK. The benefits of exercise combined with physiotherapy in the treatment of adults with cystic fibrosis. *Respiratory Medicine* 1992; 86:507-11.
  42. Dwyer TJ, Alison JA, McKeough ZJ, Daviskas E, Bye PTP. Effects of exercise on respiratory flow and sputum properties in patients with cystic fibrosis. *Chest* 2011; 139:870-7.
  43. Lee AL, Rawlings S, Bennett KA, Armstrong D. Pain and its clinical associations in individuals with cystic fibrosis: A systematic review. *Chron Respir Dis* 2016; 13:102-17.
  44. Hayes M, Yaster M, Haythornthwaite JA, et al. Pain is a common problem affecting clinical outcomes in adults with cystic fibrosis. *Chest* 2011; 140:1598-603.
  45. Barker N, Raghavan A, Buttling P, Douros K, Lloyd Everard M. Thoracic kyphosis is now uncommon amongst children and adolescents with cystic fibrosis. *Front Pediatr* 2014; 2:11.
  46. Tejero Garcia S, Giráldez Sánchez MA, Cejudo P, et al. Bone health, daily physical activity, and exercise tolerance in patients with cystic fibrosis. *Chest* 2011; 140:475-81.
  47. Sermet-Gaudelus I, Castanet M, Retsch-Bogart G, Aris MR. Update on cystic fibrosis-related bone disease: A special focus on children. *Paediatric Respiratory Reviews* 2009; 10:134-42.
  48. White D, Stiller K, Roney F. The prevalence and severity of symptoms of incontinence in adult cystic fibrosis patients. *Physiotherapy Theory and Practice* 2000; 16:35-42.
  49. Cornacchia M, Zenorini A, Perobelli S, Zanolli L, Mastella G, Braggion C. Prevalence of urinary incontinence in women with cystic fibrosis. *BJU International* 2001; 88:44-8.
  50. Moran F, Bradley JM, Boyle L, Elborn JS. Incontinence in adult females with cystic fibrosis: A northern Ireland survey. *International Journal of Clinical Practice* 2003; 57:182-3.
  51. Burge AT, Holland AE, Sherburn M, et al. Prevalence and impact of urinary incontinence in men with cystic fibrosis. *Physiotherapy* 2015; 101:166-70.
  52. Dodd ME, Prasad SA. Physiotherapy management of cystic fibrosis. *Chronic Respiratory Disease* 2005; 2:139-49.
  53. Orr A, McVean RJ, Webb AK, Dodd ME, et al. Questionnaire survey of urinary incontinence in women with cystic fibrosis. *BMJ* 2001; 322:1521.
  54. Abbott J. Health-related quality of life measurement in cystic fibrosis: advances and limitations. *Chron Respir Dis* 2009; 6:31-41.
  55. Dodd ME, Langman H. Urinary incontinence in cystic fibrosis. *Journal of the Royal Society of Medicine* 2005; 98(Suppl 45):28-36.
  56. Reichman G, De Boe V, Braeckman J, Michielsen D. Urinary incontinence in patients with cystic fibrosis. *Scandinavian Journal of Urology* 2016; 50:128-31.
  57. McVean RJ, Orr A, Webb AK, et al. Treatment of urinary incontinence in cystic fibrosis. *Journal of Cystic Fibrosis* 2003; 2:171-6.
  58. Nankivell G, Caldwell P, Follett J. Urinary incontinence in adolescent females with cystic fibrosis. *Paediatric Respiratory Reviews* 2010; 11:95-9.
  59. Moran A, Brunzell C, Cohen RC, et al. Clinical care guidelines for cystic fibrosis-related diabetes: a position statement of the American Diabetes Association and a clinical practice guideline of the Cystic Fibrosis Foundation, endorsed by the Pediatric Endocrine Society. *Diabetes Care* 2010; 33:2697-708.
  60. Kelly A, Moran A. Update on cystic fibrosis-related diabetes. *J Cyst Fibros* 2013; 12:318-31.
  61. Lannig S. Glucose intolerance in cystic fibrosis patients. *Paediatr Respir Rev* 2001; 2:253-9.
  62. Antoniou S, Elston C. Cystic fibrosis. *Medicine* 2016; 44:321-5.
  63. Abbott J, Hurley AM, Morton MA, Conway PS. Longitudinal association between lung function and health-related quality of life in cystic fibrosis. *Thorax* 2013; 68:149-54.
  64. Gee L, Abbott J, Conway S, Etherington C, Webb A. Development of a disease specific health related quality of life measure for adults and adolescents with cystic fibrosis. *Thorax* 2000; 55:946-54.
  65. Ward N, Stiller K, Rowe H, Holland AE. The psychometric properties of the Leicester Cough Questionnaire and Respiratory Symptoms in CF tool in cystic fibrosis: A preliminary study. *J Cyst Fibros* 2017; 16:425-32.
  66. Abbott J, Hart A, Havermans T, et al. Measuring health-related quality of life in clinical trials in cystic fibrosis. *J Cyst Fibros* 2011; 10(Suppl 2):S82-5.
  67. Modi AC, Quittner AL. Validation of a disease-specific measure of health-related quality of life for children with cystic fibrosis. *J Pediatr Psychol* 2003; 28:535-45.
  68. Henry B, Aussage P, Grosskopf C, Goehrs JM. Development of the Cystic Fibrosis Questionnaire (CFQ) for assessing quality of life in pediatric and adult patients. *Qual Life Res* 2003; 12:63-76.
  69. Quittner AL, Buu A, Messer MA, Modi AC, Watrous M. Development and validation of The Cystic Fibrosis Questionnaire in the United States: a health-related quality-of-life measure for cystic fibrosis. *Chest* 2005; 128:2347-54.
  70. Alpern AN, Brumback LC, Ratjen F, Rosenfeld M, Davis SD, Quittner AL. Initial evaluation of the Parent Cystic Fibrosis Questionnaire--Revised (CFQ-R) in infants and young children. *J Cyst Fibros* 2015; 14:403-11.
  71. Elphick HE, Mallory G. Oxygen therapy for cystic fibrosis. *Cochrane Database of Systematic Reviews* 2013; (7):CD003884. doi: 10.1002/14651858.CD003884.pub4.
  72. McKone EF, Barry SC, FitzGerald MX, Gallagher CG. The role of



- supplemental oxygen during submaximal exercise in patients with cystic fibrosis. *Eur Respir J* 2002; 20:134-42.
73. Hardinge M, Annandale J, Bourne S, et al. British Thoracic Society guidelines for home oxygen use in adults. *Thorax* 2015; 70(Suppl 1):i1-43.
  74. Flume PA, Robinson KA, O'Sullivan BP, et al; Clinical Practice Guidelines for Pulmonary Therapies Committee. Cystic Fibrosis Pulmonary Guidelines: Airway clearance therapies. *Respiratory Care* 2009; 54:522-37.
  75. Stanford G, Parrott H, Bilton D, Agent P. Positive pressure-analysing the effect of the addition of non-invasive ventilation (NIV) to home airway clearance techniques (ACT) in adult cystic fibrosis (CF) patients. *Physiother Theory Pract* 2015; 31:270-4.
  76. Moran F, Bradley JM, Piper AJ. Non-invasive ventilation for cystic fibrosis. *Cochrane Database of Systematic Reviews* 2013;(4):CD002769. doi: 10.1002/14651858.CD002769.pub4.
  77. Oates GR, Stepanikova I, Gamble S, Gutierrez HH, Harris WT. Adherence to airway clearance therapy in pediatric cystic fibrosis: Socioeconomic factors and respiratory outcomes. *Pediatr Pulmonol* 2015; 50:1244-52.
  78. Proesmans M. Best practices in the treatment of early cystic fibrosis lung disease. *Therapeutic Advances in Respiratory Disease* 2017; 11:97-104.
  79. Davidson KL. Airway clearance strategies for the pediatric patient. *Respir Care* 2002; 47:823-8.
  80. Button BM, Heine RG, Catto-Smith AG, Phelan PD, Olinsky A. Chest physiotherapy, gastro-oesophageal reflux, and arousal in infants with cystic fibrosis. *Arch Dis Child* 2004; 89:435-9.
  81. Santos MD, Milross MA, Eisenhuth JP, Alison JA. Pressures and oscillation frequencies generated by bubble-positive expiratory pressure devices. *Respir Care* 2017; 62:444-50.
  82. Flume PA, O'Sullivan BP, Robinson KA, et al; Cystic Fibrosis Foundation, Pulmonary Therapies Committee. Cystic fibrosis pulmonary guidelines: chronic medications for maintenance of lung health. *Am J Respir Crit Care Med* 2007; 176:957-69.
  83. Lannefors L, Button BM, McIlwaine M. Physiotherapy in infants and young children with cystic fibrosis: current practice and future developments. *J R Soc Med* 2004; 97:(Suppl 44):8-25.
  84. Rowe SM, Miller S, Sorscher EJ. Cystic fibrosis. *N Engl J Med* 2005; 352:1992-2001.
  85. Button BM, Wilson C, Dentice R, et al. Physiotherapy for cystic fibrosis in Australia and New Zealand: A clinical practice guideline. *Respirology* 2016; 21:656-67.